



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

May 29, 2018

Nathan P. Ehresman
Sr. Regulatory Scientist
Nufarm Americas Inc.-AGT Division
c/o Nufarm Americas Inc.
4020 Aerial Center Parkway
Morrisville, NC 27560

Subject: Label Amendment – Addition of directions for use optional label statements; label updates
Product Name: EQUI-TM 1.5
EPA Registration Number: 55146-154
Application Date: 07/31/2017
Decision Number: 532157

Dear Mr. Ehresman:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Page 2 of 2
EPA Reg. No. 55146-154
Decision No. 532157

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Maryam K. Muhammad by phone at 703-347-0301, or via email at Muhammad.maryam@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to be 'HJ', written over a circular scribble.

Hope Johnson, Product Manager 21
Fungicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

METALAXYL	GROUP	4	FUNGICIDE
THIOPHANAT-METHYL	GROUP	1	FUNGICIDE
FLUDIOXONIL	GROUP	12	FUNGICIDE

EQUI-TM 1.5

[ALTERNATE BRAND NAMES: SPIRATO MTM 285 FS]

For use as a seed treatment for protection of seed and seedlings against listed diseases on soybeans, dry beans, and snap beans

ACTIVE INGREDIENT:

Metalaxyl..... 14.26%
 Thiophanate -methyl 9.30%
 Fludioxonil 2.31%

OTHER INGREDIENTS: 74.13%
TOTAL: 100.00%

Contains 1.31 lbs metalaxyl, 0.86 lbs thiophanate-methyl, and 0.21 lbs fludioxonil per gallon.

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

**For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
 For Medical Emergencies Only, Call (877) 325-1840**

EPA REG. NO. 55146-154
 EPA EST. NO. _____

MANUFACTURED FOR
 NUFARM AMERICAS INC.
 AGT DIVISION
 11901 S. Austin Avenue
 Alsip, IL 60803



NET CONTENTS _____ GAL. (_____ Liters)
 [Designation as "NONREFILLABLE" or "REFILLABLE" for containers > 5 GAL]

[Grow a better tomorrow.]

ACCEPTED
 May 29, 2018
 Under the Federal Insecticide, Fungicide
 and Rodenticide Act as amended, for the
 pesticide registered under
 EPA Reg. No. 55146-154

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks,
- Chemical-resistant gloves for all mixers and loaders and for applicators using hand held equipment, and
- Chemical-resistant apron for mixers, loaders and other handlers exposed to the concentrate.

Note: Persons involved in bagging treated seed, sewing or moving bags of treated seed, or cleaning up bagging areas or seed treatment equipment are pesticide handlers and must wear the PPE required on this label for pesticide handlers.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

FIRST AID

IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything to an unconscious person.
IF INHALED	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.• Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. For terrestrial use, **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment wash water or rinsate. Cover or incorporate spilled treated seeds.

Groundwater Advisory: Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

PHYSICAL AND CHEMICAL HAZARDS

DO NOT mix or allow coming in contact with oxidizing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

EQUI-TM 1.5 is formulated for on-farm or commercial seed treatment using standard slurry or mist-type seed treatment equipment.

Mix thoroughly before use or use entire container at one time. All tank mixes should be pre-tested to determine physical compatibility between formulations. Observe the most restrictive use precautions and limitations on labeling of the products used in mixtures.

EQUI-TM 1.5 contains an EPA approved dye/colorant that imparts an unnatural color to the seed.

Treated seed must not be used for or mixed with food or animal feed or processed for oil. Seed commercially treated with this product must be labeled in accordance with all applicable requirements of the Federal Seed Act.

[NOTE TO REVIEWER: *[One of the following statements will be used on the final printed labeling. Nufarm reserves the option to choose the statement that will be used].*

[STATEMENT 1: Use the entire contents of EQUI-TM 1.5 in this container in the use season immediately following purchase from Nufarm. Store at temperatures above 40° F but below 80° F.]

[STATEMENT 2: The EQUI-TM 1.5 in this container has been manufactured for use in this use season. Use the entire contents of EQUI-TM 1.5 in this container in the use season immediately following purchase from Nufarm. Store at temperatures above 40° F but below 80° F. Remaining product is non-returnable.]

PRODUCT INFORMATION

EQUI-TM 1.5 is a seed treatment fungicide for use against damping-off, seed, and seedling rots due to *Pythium*, *Fusarium*, *Rhizoctonia* spp. In soybeans, dry beans, and snap beans. EQUI-TM 1.5 also suppresses seed-borne *Sclerotinia* and *Phomopsis* spp. particularly in soybeans.

Apply as a seed treatment using standard slurry or mist-type seed treatment equipment. Uniform application to seed is necessary to ensure seed safety and best disease protection. Seed must be sound and well cured prior to treatment. Product must be diluted with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding slurry rates for the crop to be treated with this product.

Resistance Management Recommendations: Equi-TM 1.5 contains metalaxyl, a Group 4 fungicide, thiophanate-methyl, a Group 1 fungicide, and fludioxonil, a Group 12 fungicide. Metalaxyl belongs to the phenylamide class of chemistry which interferes with fungal RNA synthesis. Thiophanate-methyl belongs to the Methyl Benzimidazole Carbamates class of chemistry which interferes with mitosis. Fludioxonil belongs to the phenylpyrrole class of chemistry which interferes with osmotic signal transduction.

Fungal populations may contain individuals naturally resistant to Group 1, Group 4 and 12 fungicides and if used repeatedly in the same fields, then resistant members may eventually dominate the population. Because resistance development cannot be predicted, the use of this product should conform to sound resistance management strategies such as an alternation with fungicides of a different mode of action and/or tank mixes established for the crop and use area.

Use should be based on an IPM program that includes field sanitation, scouting, historical information related to pesticide use, and crop rotation. The IPM program should also consider cultural, biological, and other chemical control practices.

Nufarm encourages responsible product stewardship to ensure effective long term control of the fungal diseases on this label.

For additional information on Fungicide Resistance Management:

- Contact Nufarm Americas Inc representatives at 1-800-345-3330
- Contact your local extension specialist or certified crop advisor.
- Visit the Fungicide Resistance Action Committee (FRAC) at <http://www.frac.info>

Tank Mixing: All tank mixes should be pre-tested to determine physical compatibility between formulations. When tank mixing, follow the most restrictive label instructions for any tank mix partner. DO NOT tank mix with any product that bears a label prohibition against tank mixing. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

MIXING INSTRUCTIONS FOR SOYBEANS

EQUI-TM 1.5 is formulated for on-farm or commercial treatment to be used with liquid rhizobia products, using standard mechanical slurry or mist-type seed treatment equipment which accurately measures and mixes a flowable seed treatment. The equipment must provide uniform coverage of EQUI-TM 1.5 on the seed. Uneven seed coverage may not give the desired level of disease control. EQUI-TM 1.5 must be diluted with either water or liquid inoculants prior to application to soybeans. Consult the manufacturer of the application equipment you plan to use for suitability for this application and for instructions on operation and calibration of the equipment. Do not mix EQUI-TM 1.5 fungicide with other pesticides unless sufficient compatibility, loading rates, and seed and crop safety factors are determined. Do not mix with any product that contains a prohibition on mixing. Follow the more restrictive labeling of any tank-mix partner.

In a clean container large enough to contain the entire mixture add the specified amount of EQUI-TM 1.5. Add half of the required water or liquid inoculant and mix thoroughly. Add the remaining amount of water or liquid inoculant and mix again. The slurry is ready for application to the soybeans. Mix as needed and do not store diluted mixture.

Additional or continuous agitation or mixing of the slurry mixture may be necessary to prevent the EQUI-TM 1.5 from settling out of solution when using water or liquid inoculants. EQUI-TM 1.5 has been found to be compatible with some liquid inoculant products. EQUI-TM 1.5 may be mixed or applied sequentially with approved liquid inoculants. Consult the maker of the liquid inoculants and a Nufarm Americas representative for directions before applying EQUI-TM 1.5 with liquid inoculants.

Note: EQUI-TM 1.5 is formulated to be compatible with some liquid inoculants. If water is used to further dilute EQUI-TM 1.5 when rhizobia are present, use only nonchlorinated water. Chlorinated water will kill the live rhizobia contained in the inoculant.

MIXING INSTRUCTIONS FOR OTHER BEANS (DRY and SUCCULENT)

Apply EQUI-TM 1.5 as a water based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of disease protection. Thoroughly mix the labeled amount of EQUI-TM 1.5 into the required amount of water for the slurry treater and dilution rate to be used.

Follow the manufacturer application instructions for the seed treatment equipment being used. Maintain constant agitation of the slurry during the treatment. Allow seed to dry before bagging.

USE RESTRICTIONS

DO NOT use treated seed for feed, food, or oil.

DO NOT use for hopper box or planter box seed treatments.

SEED BAG LABELING REQUIREMENTS

Seed commercially treated with this product must be labeled in accordance with all applicable requirements of the Federal Seed Act. The user of this product is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.

THE FEDERAL SEED ACT REQUIRES THAT BAGS CONTAINING TREATED SEED BE LABELED WITH THE FOLLOWING INFORMATION:

- This seed has been treated with EQUI-TM 1.5 (metalaxyl, thiophanate-methyl, fludioxonil) seed treatment.
- **DO NOT** use for food, feed, or oil purposes. Store away from feeds and other foodstuffs.

THE U.S. ENVIRONMENTAL PROTECTION AGENCY REQUIRES THE FOLLOWING STATEMENTS ON CONTAINERS CONTAINING TREATED SEED:

- Store treated seed away from feeds and foodstuffs.
- Wear long pants, long-sleeved shirt, shoes, socks, and chemical-resistant gloves when handling treated seed.
- Treated seed exposed on soil surface may be hazardous to birds and wildlife.
- Dispose of all excess treated seed by burying seed away from bodies of water. Cover or collect treated seeds spilled during loading and planting.
- **DO NOT** contaminate bodies of water when disposing of planting equipment wash water.
- Dispose of seed packaging in accordance with local requirements.
- **DO NOT** contaminate water bodies when disposing of planting equipment wash water.
- **DO NOT** allow children, pets, or livestock to have access to treated seed.
- **DO NOT** graze or feed livestock on treated areas for 30 days after planting.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.
- **Groundwater Advisory:** Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.
- After the seeds have been planted, do not enter or allow worker entry into treated areas during the **restricted-entry interval (REI) of 48 hours**. Exception: If the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, chemical-resistant gloves made of any waterproof material, and chemical-resistant footwear.

CROP SPECIFIC USE DIRECTIONS

SOYBEANS					
DISEASE	USE RATE (based on 3,000 seeds / lb)				
Protection against damping-off and seed and seedling rots caused by <i>Pythium</i> , <i>Fusarium</i> , and <i>Rhizoctonia</i> spp., and others Protection against <i>Phytophthora</i> (see remarks below) Suppresses seed-borne <i>Sclerotinia</i> and <i>Phomopsis</i> spp.	fl.oz. / cwt. seed*	fl.oz / 1,000 seed	fl.oz / 140,000 seed	mg AI / seed	
		1.5	0.005	0.7	0.0232 metalaxyl 0.0152 thiophanate-methyl 0.0037 fludioxonil
		*Contains 4.642 g of metalaxyl, 3.048 g of thiophanate-methyl, and 0.744 g of fludioxonil per fl.oz (100 kg seed contains 15.35 grams metalaxyl, 10.08 grams thiophanate-methyl, and 2.28 grams fludioxonil)			
Apply EQUI-TM 1.5 at the rate specified in the rate table, plus water to prepare a slurry of sufficient volume and viscosity to adequately cover the seed. As an alternative to water, you may use approved liquid <i>rhizobia</i> inoculants at rates specified by the manufacturer and may add the inoculants to the slurry to achieve sufficient total slurry volume and viscosity to adequately cover the seed.					
The addition of an application of SEBRING 318 FS or Sebring 480 FS or another approved metalaxyl or mefenoxam containing seed treatment product may be used if targeting fields with high <i>Phytophthora</i> pressure. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.					
RESTRICTIONS					
After the seeds have been planted, do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours . Exception: If the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, chemical-resistant gloves made of any waterproof material, and chemical-resistant footwear.					

DRY BEANS, SUCCULENT (SNAP) BEANS (<i>Cicer arietinum</i> (chickpea, garbanzo bean); <i>Lupinus</i> spp. (sweet lupine, white sweet lupine, white lupine, and grain lupine). <i>Phaseolus</i> spp. (kidney bean, lima bean, mung bean, navy bean, pinto bean, snap bean, and waxbean); <i>Vicia faba</i> (broad bean, fava bean); <i>Vigna</i> spp. (asparagus bean, blackeyed pea and cowpea))	
DISEASE	USE RATE (fl.oz / cwt. seed)
For protection against damping-off and seed and seedling rots caused by <i>Pythium</i> , <i>Fusarium</i> , and <i>Rhizoctonia</i> spp. and others. Protection against <i>Phytophthora</i> (see remarks below) Suppresses seed-borne <i>Sclerotinia</i> and <i>Phomopsis</i> spp.	1.5
The addition of an application of SEBRING 318 FS or Sebring 480 FS or another approved metalaxyl or mefenoxam containing seed treatment product may be used if targeting fields with high <i>Phytophthora</i> pressure. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.	
RESTRICTIONS	
After the seeds have been planted, do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours . Exception: If the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, chemical-resistant gloves made of any waterproof material, and chemical-resistant footwear.	

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool place, **DO NOT** store in direct sunlight. Protect from freezing temperatures.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: [[Note to Reviewer: The following statement will be included on all Final Printed Labels bearing multiple Container Handling statements] **NOTE:** This product is available in multiple containers. Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable" or "Refillable" designation. Follow the container handling instructions below that apply to your container type / size."]

[Nonrefillable Containers 5 Gallons or Less]

Nonrefillable container. DO NOT reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

[Nonrefillable containers larger than 5 gallons]

Nonrefillable container. DO NOT reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

[Refillable containers larger than 5 gallons]

Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

[Refillable containers for return to Nufarm]

Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Close all openings and replace all caps. Contact Nufarm's Customer Service Department at 1-800-345-3330 to arrange for return of the empty refillable container.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1)THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BYWAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BYTHE BUYER, USER, OR ITS CUSTOMERS.TOTHE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV052918)

Sebring and Spirato are registered trademarks of Nufarm Americas Inc.

All other trademarks are the property of their respective owners.

OPTIONAL MARKETING CLAIMS

[TO BE INCLUDED ON PRODUCT INFORMATION BULLETINS, MARKETING LITERATURE, AND COMPANY WEBSITES]

Three Powerful Actives, One Strong Start.

Protect soybeans with EQUI-TM 1.5.

The registered blend of three proven fungicides deliver powerful protection against the devastating diseases Rhizoctonia and Fusarium without the work and worries of a custom blend order.

Key Performance Benefits

Fludioxonil protects seeds against labeled rots and young seedlings against Rhizoctonia and Fusarium

The labeled rate of metalaxyl provides both Pythium and Phytophthora control

Thiophanate-methyl works with fludioxonil for increased control and systemic disease protection on listed crops.

Convenient to use formulation and packaging

[Not included in Final Printed Labeling]

LABEL HISTORY

File Name	Revision Mark	Comment
055146-00xxx.20150326.DRAFT (NUP-14052)	RV032615	Draft Label
055146-00xxx.20151218.DRAFT (NUP-14052)	RV121815	Revised as per EPA comments
055146-00RLU.20151221.DRAFT (NUP-14052)	RV122115	Additional changes EPA comments
055146-00154.20151229.MASTER	RV122915	FINAL APPROVAL
055146-00154.20160112.DRAFT	RV011216	Notification of ABN & LBS/Gallon update
055146-00154.20160201.MASTER	RV020116	EPA ACCEPTANCE
055146-00154.20160308.DRAFT	RV032816	Notification to add branding tagline & optional statement on page 3.
055146-00154.20160329.MASTER	RV032916	EPA ACCEPTANCE
055146-00154.20170123.DRAFT	RV012317	Notification to Add Optional Marketing Claims Page
055146-00154.20170210.DRAFT	RV021017	Revisions requested by EPA review
055146-00154.20170214.DRAFT	RV021417	EPA ACCEPTANCE
055146-00154.20170731.DRAFT	RV073117	Addition of Optional Statement
055146-00154.20171030.DRAFT	RV102617	Updated Statements Per EPA Reviewer
055156-00154.20180522.DRAFT	RV052218	Updated Statements Per EPA Review
055146-00154.20180529.DRAFT	RV052918	Revisions requested by EPA